

**TISSUE DOPPLER IMAGING FOR THE DIAGNOSIS OF RIGHT
VENTRICULAR MYOCARDIAL INFARCTION AND
IDENTIFICATION OF INFARCT RELATED ARTERY IN ACUTE
INFERIOR WALL MYOCARDIAL INFARCTION.**

**DISSERTATION SUBMITTED
IN PARTIAL FULFILLMENT OF THE REGULATIONS
FOR THE AWARD OF THE DEGREE OF**

DM BRANCH – II

CARDIOLOGY

STANLEY MEDICAL COLLEGE, CHENNAI

AUGUST 2009



**THE TAMILNADU Dr.M.G.R. MEDICAL UNIVERSITY
CHENNAI**

CERTIFICATE

This is to certify that the dissertation title “**TISSUE DOPPLER IMAGING FOR THE DIAGNOSIS OF RIGHT VENTRICULAR MYOCARDIAL INFARCTION AND IDENTIFICATION OF INFARCT RELATED ARTERY IN ACUTE INFERIOR WALL MYOCARDIAL INFARCTION**” is the bonafide work of
DR.P.THIRUMURUGAN in partial fulfillment of the requirements for D.M
Branch-II (cardiology) Examination of The Tamilnadu DR.M.G.R. Medical
University to be held in August 2009.The period of study was from January
2007-December 2008

PROF. R. SUBRAMANIAN M.D., D.M.,
Professor and Head of the
Dept.of Cardiology,
Govt. Stanley medical college & Hospital
Chennai-600 001.

DEAN
Govt. Stanley medical college & Hospital,
Chennai-600 001.

DECLARATION

**I Dr. P.THIRUMURUGAN., solemnly declare that the
dissertation titled**

**TISSUE DOPPLER IMAGING FOR THE DIAGNOSIS OF RIGHT
VENTRICULAR MYOCARDIAL INFARCTION AND IDENTIFICATION OF
INFARCT RELATED ARTERY IN ACUTE INFERIOR WALL MYOCARDIAL
INFARCTION is a bonafide work done by me at Govt.Stanley Medical College and
Hospital during January 2007 to December 2008 under the guidance and
supervision of my PROF.DR. R.SUBRAMANIAN,M.D.,D.M Professor and Head of
Dept.of Cardiology.**

**This dissertation is submitted to The Tamilnadu DR. M.G.R Medical
University, towards partial fulfillment of requirement for the award of D.M Degree
Branch-II Cardiology**

**Chennai
Date:**

Place:

(Dr. P. THIRUMURUGAN)

ACKNOWLEDGEMENT

I owe my thanks to the Dean, Govt.Stanley Medical College and Hospital, DR.J. MOHANASUNDRAM M.D, PhD, DNB. for me to avail the facilities needed for my dissertation work.

I wish to express my respect and sincere gratitude to my beloved teacher PROF. DR. R. SUBRAMANIAN, M.D., D.M (CARDIOLOGY) Professor and Head of Department of Cardiology for his valuable guidance and encouragement through out the study.

I am extremely thankful to our PROF. DR. M. SOMASUNDRAM, M.D., D.M, (CARDIOLOGY) Additional Professor of Cardiology for his support and guidance during the study.

I am also expressing my gratitude to my assistant professors DR. DR.K.KANNAN., DR. M. NANDAKUMARAN., DR. SIVAN., DR.ASHOKVICTOR, DR.P.M.NAGESWARAN, DR.K.TAMILSELVAN, DR. GUNASEKARAN., DR. S. JEYABOOPATHY.,DR.A.RAVICHANDRAN

Last but not the least, my sincere thanks to all the patients who cooperated for the study. ,

CONTENTS

1. INTRODUCTION	:	1
2. REVIEW OF LITERATURE	:	23
3. AIM OF STUDY	:	34
4. MATERIALS AND METHODS	:	35
5. RESULTS	:	39
6. DISCUSSION	:	46
7. LIMITATIONS	:	50
8. CONCLUSION	:	51
9. BIBLIOGRAPHY	:	52

INTRODUCTION

In approximately half of patients with inferior wall myocardial infarction, right ventricular myocardial infarction (RVMI) develops. In patients with RVMI, the risk of death in the hospital is high and major complications are greater. For early diagnosis, electrocardiography and two dimensional echocardiography are used, but these methods are occasionally insufficient.

Tissue Doppler imaging (TDI) has evolved as a new technique that enables myocardial velocities to be detected and makes the quantitative assessment of the systolic and diastolic movements of myocardial walls possible.

This study was designed to test the usefulness both of the right ventricular peak myocardial systolic velocity (S_m) and of the myocardial performance index (MPI) obtained by pulsed –wave TDI in the diagnosis of RVMI and in determining the infarct-related artery (IRA) in patients with acute inferior wall myocardial infarctions.

RIGHT VENTRICULAR INFARCTION

Frequency

Right ventricular infarction usually is noted in association with inferior wall myocardial infarction. The incidence of right ventricular infarction in such cases ranges from 10-50%, depending on the series (Andersen, Nielsen, Falk et al, 1989). Isolated infarction of the right ventricle is extremely rare.

Although right ventricular infarction is clinically evident in a sizable number of cases, the incidence is considerably less than that found at autopsy (Andersen, 1987; Andersen, Nielsen, Lund, 1989; Bates 1997; Birnbaum et al, 1999). A major reason for the discrepancy is the difficulty in establishing the presence of right ventricular infarction in living subjects. Additionally, right ventricular dysfunction and stunning frequently is of a transient nature, such that estimation of its true incidence is even more difficult. Criteria have been set for to diagnose right ventricular infarction; even when strictly employed, however, the criteria lead to underestimation of the true incidence of right ventricular infarction (Baat et al, 1984; Baat et al, 1983; Elkayam et al, 1979).

Importance of identifying RVM

Despite the initial observation of serious hemodynamic consequences of right ventricular infarction nearly two decades ago, this condition has received little clinical attention until recent years. Right ventricular infarction contributes markedly to hemodynamic instability, atrioventricular conduction block, and in-hospital mortality in patients with inferior myocardial infarctions.

As compared with all clinical variables available at the time of admission, right ventricular infarction was associated with a relative risk of in-hospital mortality of 7.7 (95 percent confidence interval, 2.4 to 9). Ultimately, 95 percent of patients

without evidence of right ventricular infarction at the time of admission were discharged from the hospital, as compared with only 69 percent of those in whom right ventricular infarction complicated the acute inferior myocardial infarction.

Interest in recognizing right ventricular infarction noninvasively has grown because of the therapeutic implications of distinguishing patients with right ventricular dysfunction from those without right ventricular dysfunction.

The potential hemodynamic derangements associated with right ventricular infarction render the afflicted patient unusually sensitive to diminished preload (i.e., volume) and loss of atrioventricular synchrony. These two circumstances can result in a severe decrease in right and, secondarily, left ventricular output (Hirsowitz et al, 1984; Hurst, 1998; Iqbal and Liebson, 1981).

Infarct Related artery

The posterior descending branch of the right coronary artery usually supplies the inferior and posterior walls of the right ventricle. The marginal branches of the right coronary artery supply the lateral wall of the right ventricle. The anterior wall of the right ventricle has a dual blood supply: the conus branch of the right coronary artery and the moderator branch artery, which courses from the left anterior descending artery (Forman et al, 1984).

A direct correlation exists between the anatomic site of right coronary artery occlusion and the extent of right ventricular infarction. Studies have demonstrated that more proximal right coronary artery occlusions result in larger right ventricular infarctions (Garty et al, 1984). If occlusion occurs before the right ventricular marginal branches, and collateral blood flow from the left anterior descending coronary artery is absent, then the size of infarction generally is greater. Extent of infarction depends somewhat on flow through the thebesian veins (Kinn et al 1995).

In general, any major reduction in blood supply to the right ventricular free wall portends an adverse prognosis in association with this disorder⁷.

Clinical features

- ❖ A right ventricular infarct should be considered in all patients who present with an acute inferior wall myocardial infarction, especially in the setting of a low cardiac output.
- ❖ Patients may describe symptoms consistent with hypotension.
- ❖ A subtle clue to the presence of hemodynamically significant right ventricular infarction is a marked sensitivity to preload reducing agents such as nitrates, morphine, or diuretics (Mittal, 1994).
- ❖ Other presentations include high-grade atrioventricular block, tricuspid regurgitation, cardiogenic shock, right ventricular free wall rupture, and cardiac tamponade.
- ❖ Patients with extensive right ventricular necrosis are at risk for right ventricular catheter-related perforation, and passage of a floating balloon catheter or pacemaker must always be performed with great care in such a setting.

Physical Signs

- The classic clinical triad of right ventricular infarction includes distended neck veins, clear lung fields, and hypotension (Mavric et al, 1990)
- Infrequent clinical manifestations include right ventricular third and fourth heart sounds, which are typically audible at the left lower sternal border and increase with inspiration.

- On hemodynamic monitoring, disproportionate elevation of rights-sided filling pressures compared with left-sided hemodynamics represents the hallmark of right ventricular infarction.

DIAGNOSIS

Electrocardiography

- All patients with interior wall myocardial infarction should have a right-sided ECG. ST-segment elevation in lead V_{4R} predicts right ventricular involvement, identifying a high-risk subset of patients in the setting of inferior wall myocardial infarction (Robalino et al, 1990) ST- segment elevation in lead V_{4R} had an overall sensitivity of 76 percent, a specificity of 78 percent, and a diagnostic accuracy of 72 percent for the diagnosis.

Limitations of the Electrocardiogram in the Diagnosis of Right Ventricular Infarction

The use of ST-segment elevation in lead V_{4R} for the diagnosis of right ventricular infarction has its own limitations. The ST-segment elevation is transient, disappearing in less than 10 hours following its onset in half of patients. The use of ST-segment elevation in lead V_{4R} for the diagnosis of right ventricular infarction is known to lose specificity in the presence of any heart disease that may induce ST-segment elevation in lead V₁, such as pericardial disease, acute pulmonary embolism, left anterior fascicular block, and acute anterior myocardial infarction (present in 10 percent of patients with right ventricular involvement).

Imaging Studies

- In the appropriate clinical setting, a diagnosis of right ventricular infarction can be made using noninvasive techniques, or the patients may require right ventricular catheterization and hemodynamic monitoring.

Echocardiography

- Right ventricular dilatation, abnormal right ventricular wall motion, paradoxical motion of the interventricular septum and tricuspid regurgitation are echocardiographic features of right ventricular infarction.
- It is difficult to assess both right ventricular anatomy and function with reliable conventional echocardiographic examination because of its complex structure. In most cases, the visual assessment of the right ventricular free wall by echocardiography leads to an underestimation of hypokinesia, due to an asymmetric contraction of the right ventricular walls toward its center.
- In the vast majority of patients with right ventricular infarction, the wall motion abnormalities initially manifest on echocardiography reverse within 3 months (Strauss et al, 1980)
- Echocardiography is useful as a modality to rule out pericardial disease and tamponade, which are the major differential diagnoses in the setting of a right ventricular infarction.
- Echocardiogram has 82% sensitivity and 93% specificity in detecting right ventricular infarction when right ventricular scintigraphy is used as the comparative standard (Singhal et al, 1984).
- Gated equilibrium radio nuclide angiography and technetium 99m pyrophosphate scintigraphy are useful in diagnosing right ventricular infarction noninvasively (Sugimoto et al, 1996). In the case of radionuclide angiography, the right ventricle is demonstrated to be enlarged and poorly contractile, with a reduced ejection fraction. When technetium 99m pyrophosphate is employed, the right ventricular free wall is "hot", indicating

significant infarction.

TISSUE DOPPLER ECHOCARDIOGRAPHY

Tissue Doppler (TD) imaging is a novel echocardiographic technique that measures myocardial velocities. However, there are sparse data on TD imaging of the right ventricular (RV) free wall in the diagnosis and prognosis of RV myocardial infarction (M1) in inferior wall left ventricular M1. As determined from the position of the right ventricle tricuspid annulus by TDI, myocardial velocities and the myocardial performance index (MPI) can give information about right ventricular function. The evaluation of velocities of using DTI provides a noninvasive and rapid method for assessing right ventricular function in patients with RVMI.

HEMODYNAMIC MONITORING

Disproportionate elevation of right-sided filling pressures when compared with left-sided hemodynamics represents the hallmark of right ventricular infarction.

Accepted hemodynamic criteria for right ventricular infarction include right atrial pressure greater than 10 mm Hg, right atrial –to-pulmonary capillary wedge pressure ratio greater than 0.8. These values may manifest only after volume loading (Sugiura et al, 1994). Other interesting hemodynamic features of right ventricular infarction include the following:

- Prominent "y" descent of the right atrial pressure.
- Increase in venous or right atrial pressure with inspiration (i.e., Kussmaul sign)

TISSUE DOPPLER IMAGING (TDI)

The pulsed and color Doppler echocardiography can detect motion and velocity of both moving blood and myocardial tissue. We have traditionally processed

blood flow signals in the conventional Doppler flow imaging, so that the high velocity and low amplitude Doppler signals reflected by the moving blood are measured whereas the myocardial Doppler signals with high amplitude but low velocity are suppressed.

TDI is a new ultrasound technique that is based on color Doppler imaging principles and allows quantification of intramural myocardial velocities by detection of consecutive phase shifts of the ultrasound signal reflected from the contracting myocardium. To display regional myocardial velocities, thresholding and filtering algorithms are changed to reject the low-amplitude echoes from the blood pool. These improve the ability to measure low velocity myocardial signals that are typically in the range of 0.6 – 24 cm/s.

PRINCIPLES OF TISSUE DOPPLER IMAGING

Unlike conventional Doppler signals that are typified by high velocity and low amplitude, myocardial motion is characterized by relatively low velocity and high amplitude signals. Tissue motion creates Doppler Shifts that are approximately 40 dB higher than Doppler signals from blood flow whereas velocities rarely exceed 20cm/s. To record low wall motion velocity, gain amplification is reduced and high pass filters are bypassed with the tissue signal directly entered into the autocorrelator. During image acquisition, it is important to optimise the frame rate using an image sector as narrow as possible and to select the appropriate velocity scale. These parameters should be optimised at the time of imaging, as it is not possible to modify the frame rate and the velocity scale during post processing image analysis.

MODALITIES OF TISSUE DOPPLER ECHOCARDIOGRAPHY

TDE has three modalities: spectral pulsed wave Doppler, two dimensional, and M mode colour Doppler.

Spectral Pulsed Doppler

Spectral pulsed TDI has the advantage of online measurements of velocities and time intervals and an excellent temporal resolution (8ms). According to the Doppler principle, tissue velocities moving toward the transducer are positive, whereas velocities moving away from the transducer are negative. The spectral PW-TDI method provides higher temporal resolution and resolves all peak velocities. With this modality a sample volume is placed within the myocardium (either in the endocardium or the epicardium) and the low Doppler shift of frequencies recorded from the heart wall moving through the sample volume during the cardiac cycle is recorded.

The pattern can be divided into two parts systolic and diastolic, from which several measurements can be obtained :

1. The systolic phase is characterized by a positive wave (S) preceded by the time taken for regional isovolumic contraction (IVCT);
2. The diastolic phase, which is complex, is composed of 4 periods:
 - a) Regional isovolumic relaxation (IVRT);
 - b) The rapid filling period characterized by a negative wave (E);
 - c) Diastasis, and
 - d) Filling due to atrial contraction, represented by a second negative wave(A)

COLOUR DOPPLER

In colour TDI, red encodes wall motion towards the transducer (positive velocities), whereas blue encodes wall motion away from the transducer (negative velocities). On each side of the scale, the brightest shades correspond to the highest velocities. Colour images require digital acquisition and storage for off-line post processing analysis. In contrast to spectral Doppler, endocardial and epicardial layers can be separately analysed. Peak and mean velocities, time velocity integral, and regional time intervals can be measured in each myocardial segment, in each myocardial layer, and in each phase of the cardiac cycle. M mode colour encoded TDI has a high temporal resolution (5-10 ms). Colour two dimensional imaging has been limited by a slow frame rate, but parallel processing and advances in beam formation technology have increased the frame rate to a level adequate for analysis of most cardiac events (temporal resolution 10-100ms).

POTENTIAL CLINICAL IMPLICATIONS OF PULSED TDI

Global assessment of LV function

The global LV systolic function can be assessed by peak mitral annular (MA) velocity in apical views. Although myocardial synchronous contraction and relaxation depends critically on coordination of circumferential and longitudinal fibers, the longitudinal fibers are particularly vulnerable to myocardial ischemia and activation disturbance likely related to their subendocardial position. Because the epicardial apex is relatively fixed, the longitudinal fiber shortening draws the atrioventricular ring towards the apex of the ventricle. Not only does this contribute to the fall in the left ventricular cavity volume with ejection, but at the same time increase the volume of the left atrium. Similarly, during diastole, the backward movement of the atrioventricular ring aids the LV filling. Dumensil et al. showed that with a 40% decrease of LVEF (M-mode) in disease, there might be only 15-20% corresponding decrease in the circumferential fiber (short-axis) shortening, highlighting the significant contribution of longitudinal fibers in overall LV function. Since the

introduction of the concept of mitral annular descent by echocardiography in 1967, it has been shown that the peak systolic mitral annular amplitude and total excursion correlate well with LV stroke volume.

Its amplitude and velocity have been used as an index of ejection fraction (EF) and been shown to correlate with the LVEF obtained by radionuclide ventriculography in humans. Tissue Doppler not only defines and measures the mitral annular movement, but also gives high temporal relationship. Decrease in peak systolic descent velocity at a mitral annular site with corresponding longer time to peak correlates with LV asynergy and infarct regions in patients with previous myocardial infarction.

The mitral annular velocity has also been shown to be a sensitive marker of alteration of LV contractility induced by inotropic stimulation. In studying 12 normal volunteers, Gorcsan et al. showed that peak mitral annular velocity increased significantly with corresponding decrease in time to peak velocity at very low dose (1mcg/kg/min) dobutamine.

A further linear dose-dependent incremental increase of the velocity was observed up to the target infusion of 5mcg/kg/min. In contrast, regional peak velocities did not change until the 2mcg/kg/min dose and no change of myocardial wall thickening and ejection fraction was observed until 3mcg/kg/min. These data support the mitral annular velocity as a sensitive marker of global systolic function, even when the endocardial definition is suboptimal for estimation of LV volumes and ejection fractions. The MA velocity provides important insight in terms of amplitude, velocity and timing in relation to the cardiac cycle.

Diastolic Function

Analysis of the motion of the myocardium during diastole with TDI has been performed in healthy normal subjects. The PW-TDI provides a mirror image of the initial inflow patterns, with a decrease in the E/A velocity ratio with age. Myocardial diastolic velocities have been reported in various diseases, allowing a differentiation

between constrictive pericarditis and restrictive cardiomyopathy, as well as among the different patterns of myocardial hypertrophy.

Mitral annular E velocities have been proposed to be preload-independent variables of diastolic function. However, subsequent works by other investigators have debated the load independence of Ea. In day-to-day practice it is often necessary to obtain a noninvasive judgment regarding the left ventricular filling pressures for guiding the management of patients with cardiac diseases. Nagueh et al. proposed a ratio of E/Ea as an important tool for noninvasive assessment of the left ventricular filling pressure.

They studied 125 patients and divided them into three groups: asymptomatic persons with left ventricular $E/A > 1$ (normal), $E/A < 1$ (impaired relaxation) and $E/A > 1$ but with symptoms of heart failure (pseudo normal). Ea was reduced in patients with impaired relaxation and pseudo normal pattern. The best correlation for pulmonary artery wedge pressure was with E/Ea ratio ($r=0.87$).

Nagueh et al. in their second study showed that the ratio of E/Ea could be used to estimate pulmonary artery wedge pressure accurately even in the presence of sinus tachycardia. They studied 100 cases, which included 35 patients with completely merging E and A mitral inflow velocity waves, and found strongest correlation of pulmonary capillary wedge pressures [PCWP] with E/Ea ratio ($r=0.86$, $PCWP=1.55+1.47 E/Ea$). The best cut-off was a value of $E/Ea > 10$ that predicted a PCWP of >15 with a 92% sensitivity and 80% specificity. Dagdelen et al. evaluated the utility of deceleration time of mitral annular Ea, Aa time and Ea/Aa ratio in estimating left ventricular end – diastolic pressure in 70 hypertensive patients with no significant coronary artery disease and normal left ventricular systolic function but with varying left ventricular end-diastolic pressures. The best sensitivity and

specificity values for predicting left ventricular end-diastolic pressure of >15 mmHg were Ea DT >120 ms (88% and 81%), Aa time 100 ms (71% and 80%) and Ea/Aa <0.5 (86% and 72%), respectively

Tissue Doppler Imaging in Coronary Artery Disease

Myocardial ischemia alters regional contraction and relaxation. An acutely ischemic myocardium suffers not only from the loss of contractility but also tensile strength and stiffness. These alterations can be quantified by tissue velocity imaging.

Derumeaux et al. studied the patterns of tissue velocity profile during graded ischemia in an animal model and observed that ischemia resulted in rapid reduction of systolic velocities and early diastolic velocities. Within 5 of occlusion of the left anterior descending (LAD) artery, a 46% reduction in systolic velocities was seen.

Systolic waves became negative at around 30s and peaked at 1 min. This corresponded to paradoxical expansion of the same segment. Diastolic abnormalities also occurred simultaneously and were characterized by a decrease in Em wave and increase in Am wave. The velocity of isometric relaxation and contraction increased and peaked at 1 min. There was a good correlation between decrease of systolic velocity and regional myocardial flow.

In another animal model, Pislaru et al. measured the Tran mural extent of necrosis in the myocardium after occlusion and relaxation velocities. The isovolumic contraction wave was found to be absent in the presence of $>20\%$ necrosis of the myocardial thickness and preserved in animals with less than 20% necrosis of thickness of the myocardium. Wall thickening did not improve with reperfusion. However, positive isovolumic contraction velocities were able to distinguish segments with different degrees of necrosis.

Preserved isovolumic contraction waves in hypokinetic or akinetic segments were associated with smaller infarcts. However, it needs to be pointed out that ischemia and perfusion-induced contractile dysfunction of myocardial perfusion through myocardial velocities, therefore, is only valid in the situation of ongoing ischemia and not following reperfusion.

In patients with chronic stable angina and normal ejection fraction, minor abnormalities of LV isovolumic relaxation and contraction and longitudinal shortening can be detected on tissue Doppler imaging (TDI). TDI has also been used during stress echocardiography for quantifying regional wall motion abnormalities. Katz et al. evaluated the utility of TDI in 60 patients for detection wall motion abnormalities. Excluding the apical segments, a peak velocity of less than 5.5 cm/s with peak stress had an average sensitivity, specificity and accuracy of 96%, 81% and 86%, respectively, for identifying abnormal segments at peak stress.

Dobutamine stress echocardiography using TDI is also an accurate tool for assessing myocardial viability. Larrazet et al. evaluated the concordance between dobutamine stress echocardiography using TDI, dobutamine stress using standard echocardiography and thallium-201 SPECT for viability assessment in 23 patients with coronary artery disease (CAD) and resting LV dysfunction. Compared to dobutamine stress using standard echocardiographic evaluation of regional wall motion abnormalities, dobutamine TDI had a higher sensitivity (60% v.82%), equal specificity (100% each) and higher accuracy (74% v.94%) for detecting viable myocardium using thallium –201 as the gold standard. Nishino et, al evaluated conventional dobutamine stress echocardiography (DSE) and color TDI (M-mode) for predicting viable myocardium in 56 patients undergoing coronary angioplasty.

Dobutamine TDI, either used alone or in combination with conventional stress

echocardiography, had a higher sensitivity and equivalent specificity for the prediction of recovery of dysfunctional segments. Tissue Doppler imaging is thus a useful technique during stress echocardiography interpretation by novices as well as experienced echocardiographers not formally trained in stress echocardiography. The tissue velocity profile is particularly important in interpreting regional wall motion in the basal segments. In a study of 77 patients who underwent dobutamine stress echocardiography and coronary angiography, TDI improved the baseline accuracy for novice readers from 68% to 76% ($p=0.001$) for basal segments, which were otherwise scored as normal.

Tissue Doppler Imaging in Right ventricular myocardial infarction

Right ventricular myocardial infarction impairs right ventricular function. However, it is difficult to assess both right ventricular anatomy and function with reliable conventional echocardiographic examination because of its complex structure. In most cases, the visual assessment of the right ventricular free wall by echocardiography leads to an underestimation of hypokinesia, due to an asymmetric contraction of the right ventricular walls toward its center. It has been suggested that color or pulsed wave TDI, which is frequently used in assessing left ventricular function, also can be used to assess right ventricular function. The myocardial velocities obtained by pulsed-wave TDI from the right ventricular free wall at the level of the tricuspid annulus in the apical four-chamber view show the longitudinal dynamics of the right ventricle, which is not seen with visual scoring.

In studies assessing right ventricular function by using techniques other than TDI, the movement of the tricuspid annulus has been shown to represent the global right ventricular function. It has been documented that significant right coronary artery disease can be identified with the assessment of systolic velocity obtained by

pulsed-wave TDI from the right ventricular free wall close to the tricuspid lateral annulus in the apical four-chamber view during dobutamine stress echocardiography. Finally, it has been shown that tricuspid annulus systolic velocity decreases in patients with RVMI complicating inferior wall MI than in controls even when there was no ECG evidence of RVMI and also in those with RVMI compared to those without RVMI.

REVIEW OF LITERATURES

In approximately half of patients with inferior wall myocardial infarction, right ventricular myocardial infarction (RVMI) develops. In patients with RVMI, the risk of death in the hospital is high and major complications are greater. Despite the initial observation of serious hemodynamic consequences of right ventricular infarction nearly two decades ago, this condition has received little clinical attention until recent years.

Right ventricular infarction contributes markedly to hemodynamic instability, atrioventricular conduction block, and in-hospital mortality in patients with inferior myocardial infarction. The following studies were conducted to evaluate RVMI and its complications.

Garty.I et al, Sixty patients with AMI were prospectively studied: 40 patients with inferior wall MI and 20 patients with anterior wall MI. Of the 40 patients diagnosed as having inferior wall MI, 20 cases (50%) were found to be associated with RVMI. 17 (85%) also demonstrated a ST segment elevation of 0.1 mV, and pathological Q waves in the V4R lead. The ejection fraction (EF) of RV was found to be significantly decreased in patients with RVMI compared with the other group (mean, 27% versus 57%). Among the 20 patients with RVMI, 16 (80%) showed various complication during the hospitalization period, versus 9 patients (45%) from the group with inferior wall MI. The most common complication in RVMI patients was conduction disturbances (7 of 20 versus 2 of 20 patients). The clinical and prognostic importance of the

early diagnosis of RVMI is stressed.

Rodrigues et al the values of several non-invasive methods for the diagnosis of right ventricular necrosis in inferior myocardial infarction were compared in 51 consecutive patients who underwent serial ECG, radionuclide ventriculography, pyrophosphate scintigraphy, and cross sectional echocardiography. Functionally important right ventricular infarction was best detected and followed serially by radionuclide ventriculography. Echocardiography methods for evaluating right ventricular ejection fraction correlated poorly with radionuclide methods.

Increased uptake of radioactivity by the right ventricle on pyrophosphate scintigraphy usually indicated poor right ventricular function, but a scan that was negative in the right ventricular territory did not exclude dysfunction. ST segment elevation in V4R was not specific for right ventricular infarction and its routine use may lead to over diagnosis of this condition. Serial measurements suggest that profound right ventricular dysfunction persists after acute inferior infarction and is associated with considerable morbidity and mortality.

Electrocardiography

All patients with inferior wall myocardial infarction should have a right-sided ECG. ST-segment elevation in lead V_{4R} predicts right ventricular involvement, identifying a high-risk subset of patients in the setting of inferior wall myocardial infarction (Robalino et al, 1990). The use of ST-segment elevation in lead V_{4R} for the diagnosis of right ventricular infarction has its own limitations. The ST-segment elevation is transient, disappearing in less than 10 hours following its onset in half of patients. The following study was conducted regarding the use of ECG in RVMI.

Kabakci.G et al this study investigated the predictive value of presentation and 24-hour electrocardiograms in defining the infarct-related artery (IRA), its lesion segment, and the right ventricular involvement in acute inferior myocardial infarction (MI). One hundred forty-nine patients with acute inferior MI were included. Infarct-related artery, its lesion segment, and the validity of new ECG criteria for the diagnosis of right ventricular MI (RVMI) were investigated. The width of Q wave in lead III > lead II criterion supported the RCA to be IRA with a sensitivity of 60% and a specificity of 61% The finding of ST elevation in V(I) but no ST elevation in (V2) on admission ECG had a sensitivity of 63% and a specificity of 99% in the diagnosis of RVMI. This study concluded that, despite high specificity, due to moderate degree

sensitivity, 12-lead ECG for the diagnosis of RVMI is questionable.

Tissue Doppler Imaging in Right ventricular myocardial infarction

It has been suggested that color or pulsed-wave TDI, which is frequently used in assessing left ventricular function, also can be used to assess right ventricular function. The myocardial velocities obtained by pulsed-wave TDI from the right ventricular free wall at the level of the tricuspid annulus in the apical four-chamber view show the longitudinal dynamics of the right ventricle, which is not seen with visual scoring.

In studies assessing right ventricular function by using techniques other than TDI, the movement of the tricuspid annulus has been shown to represent the global right ventricular function. It has been documented that significant right coronary artery disease can be identified with the assessment of systolic velocity obtained by pulsed-wave TDI from the right ventricular free wall close to the tricuspid lateral annulus in the apical four-chamber view during dobutamine stress echocardiography. The following studies were conducted to assess the use of Tissue Doppler imaging in inferior/right ventricular infarction

Pramod jaiswaal et al right ventricular (RV) function has not been studied widely after a myocardial infarction (MI). The current study was designed to determine the RV function by tricuspid annular motion and tricuspid annular velocity after MI. Ten patients with a first episode of

acute MI (inferior MI: 6, anterior MI:4) and 4 healthy controls were included in the study.

The tricuspid annular motion was reduced in inferior MI compared to controls (19.2 and 24 mm). The peak systolic velocity of tricuspid annulus was reduced in inferior MI compared to controls (12v 14.2 cm/s), also the tricuspid annular motion was significantly lower in patients with RV infarction than in those without RV infarction (16.8 and 21.6 mm, $p < 0.01$), Patients with RV infarction also had significantly decreased peak tricuspid annular systolic and early diastolic velocities. Thus, tricuspid annular motion and velocity can be used to assess RV function in association with inferior wall myocardial infarction.

Dokainish.H et al fifty patients who had left ventricular MI underwent TD echocardiography and angiography within 48 hours of MI. For diagnosis, the ability of RV TD imaging to detect RV MI was assessed using coronary angiography as the reference standard. For prognosis, the ability of TD detection of RV dysfunction to predict cardiac death or rehospitalization at 1 year was determined. On multivariate analysis, systolic annular velocity and RV dimension predicted RV MI. Decreased RV systolic annular velocity on TD images detects RV MI in first left ventricular acute inferior Mi and predicted cardiac death or rehospitalization or rehospitalization at 1 year.

Yang.S.S. et al with tissue Doppler imaging and right ventricular

Tei index, right ventricular function in patients with right ventricular myocardial infarction (RVMI) was assessed. 51 patients admitted to coronary care units and diagnosed as acute inferior myocardial infarction were studied 23 patients were diagnosed to have RVMI and 28 patients no. 20 healthy subjects served as controls. Sm and Em at the lateral side of tricuspid annulus and the RV free mid-wall reduced significantly in patients with RVMI as compared with those without RVMI and healthy individuals (Sm at the lateral (7.0 ± 2.0) cm/s vs (8.7 ± 1.9) cm/s and (10.6 ± 2.1) cm/s, $P < 0.01$; Em at the lateral (6.3 ± 1.9) cm/s vs (7.9 ± 1.8) cm/s and (9.6 ± 1.9) cm/s, $P < 0.01$; Sm at the RV free mid-wall (6.4 ± 1.9) cm/s vs (8.0 ± 1.9) cm/s and (9.4 ± 2.0) cm/s, $P < 0.05$; Em at the RV free mid-wall (6.1 ± 2.0) cm/s vs (7.6 ± 2.0) cm/s and (9.2 ± 2.3) cm/s, $P < 0.05$). RV Tei index in patients with RVMI also increased as compared with that in the other two groups (0.65 ± 0.19) vs (0.40 ± 0.15) and (0.26 ± 0.10) ; $P < 0.01$: The evaluation of velocities at the lateral side of tricuspid annulus and the RV free mid-wall using DTI and RV Tei index provides a noninvasive and rapid method for assessing right ventricular function in patients with RVMI.

Oguzhaan.A et al this study was undertaken to determine right ventricular (RV) function as assessed by colour Doppler tissue imaging (DTI) in patients with RV infarction. 35 patients were evaluated: 14 patients had an inferior myocardial infarction (MI) with RV infarction and 21 patients had an inferior MI without RV involvement. Twenty age-

matched healthy subjects served as control. Systolic and early diastolic velocities at the lateral tricuspid annulus were significantly reduced in patients with inferior MI with RV infarction compared with those in healthy individuals (7.8 ± 1 vs. 11 ± 2 cm/s, $P < 0.002$). And patients with inferior MI without RV infarction (7.8 ± 1 vs. $10 \pm$ cm/s, $P < 0.002$). The late diastolic lateral annular velocity did not differ between the groups. Systolic and early diastolic RV free wall velocities were also significantly decreased in patients with RV infarction compared with those in healthy individuals (7 ± 1 vs. 8.7 ± 1 cm/s, $P < 0.01$; 6.3 ± 2 vs. 8.7 ± 2 cm/s, $P < 0.05$, respectively and patients with inferior MI without RV infarction (7 ± 1 vs. 9 ± 2 cm/s, $p < 0.01$; 6.3 ± 2 vs. 8.3 ± 2 cm/s, $p < 0.05$, respectively). The evaluation of tricuspid annular and RV free wall velocities using colour DTI provides a rapid and noninvasive tool for assessing RV function in patients with RV infarction.

Gin.K et al this study assessed echocardiographic variables including TDI of the tricuspid valve annulus to detect RVI in the setting of acute inferior wall myocardial infarction (IWMI). 50 patients were included in the study. 22 patients (44%) had the ICL proximal to the RV branch (Group I RVMI), while 28 patients had the ICL distal to the first RV branch of the RCA or in the artery other than the RCA (Group 2 NO RVMI). The univariate predictors of RVMI were: elevation of the jugular venous pressure more than of RVMI were: elevation of the jugular

venous pressure more than of RVMI were: elevation of the jugular venous pressure more than 4cm, ST elevation in lead V4R, abnormal echo RV function by visual inspection, right ventricular dimension(RVD), and tricuspid annular systolic velocity(TAV). On logistic regression analysis including all significant univariate variables, only TAV(odds ratio 0.62, $p=0.03$) and RVD (odds ratio 1.28, $p=0.013$) predicted RVMI.

Of various clinical and echocardiographic variables, reduced systolic velocity of the tricuspid annulus by TDI and increased right ventricular dimension are multivariate predictors of RVI in patients presenting with first ST elevation IWMI.

TISSUE DOPPLER IMAGING FOR THE
DIAGNOSIS OF RIGHT VENTRICULAR
MYOCARDIAL INFARCTION AND
IDENTIFICATION OF INFARCT
RELATED ARTERY IN ACUTE INFERIOR
WALL MYOCARDIAL INFARCTION

ABSTRACT

Aim:

The aim of the study is to find out the usefulness both of peak myocardial systolic velocity (Sm) and of the myocardial performance index (MPI) of the right ventricle measured by pulsed –wave tissue Doppler imaging (TDI) to diagnose right ventricular myocardial infarction in the presence of acute inferior wall myocardial infarction.

Methods:

Fifty patients who experienced a first acute inferior MI (mean [\pm SD] age, 56 ± 8 years) were prospectively assessed. An ST-segment elevation of 0.1 mV in V4-V6R lead derivation was defined as an RVMI. From the echocardiographic apical four-chamber view, the myocardial systolic velocity Sm, the peak early diastolic velocity Em, peak late diastolic velocity Am, the ejection time, the isovolumetric relaxation time, and the contraction time of the right ventricle were recorded at the level of the tricuspid annulus by using TDI. Then, the MPI was calculated. The patients were classified into the following three groups, according to the localization of the infarct-related (IRA) detected using coronary angiography : group I, proximal right coronary artery; group II, distal right coronary artery; and group III, circumflex coronary artery.

Results:

RVMI were detected in fifteen patients, and the IRA in 21 patients was the proximal right coronary artery. The right ventricular free wall Sm was observed to be significantly low in patients with RVMI and those in group I compared to those without RVMI and those in groups II and III ($10.8 \pm 1.2 \text{ cm/s}$ vs $14.2 \pm 3.1 \text{ cm/s}$, respectively. The right ventricular free wall Sm of group I, II and III was $11.3 \pm 2.5 \text{ cm/s}$, $15.0 \pm 2.9 \text{ cm/sec}$ and $14.6 \pm 2.5 \text{ cm/s}$, respectively [$p < 0.001$]). The MPI was high in the same patient groups (0.83 ± 0.12 vs 0.57 ± 0.11) in those patients without RVMI respectively. The MPI of group I II and III was 0.74 ± 0.13 , 0.56 ± 0.15 and 0.54 ± 0.07 respectively [$p < 0.001$]). In the diagnosis of RVMI, and in the diagnosis of the proximal right coronary artery as the IRA, the sensitivity, specificity, negative predictive value, and positive predictive value of an Sm $< 12 \text{ cm/s}$ were calculated as 76%, 93%, 84%, and 86%. The sensitivity, specificity, negative predictive value, and positive predictive value of an MPI of > 0.70 in the diagnosis of RVMI were 88%, 81%, 96%, and 72%.

Conclusion:

An Sm $< 12 \text{ cm/s}$ and an MPI > 0.70 obtained by TDI may define RVMI concomitant with acute inferior myocardial infarction, and the Infarct related artery.

AIM

By using of Pulsed tissue Doppler and Myocardial performance index in assessing Right ventricular function for the diagnosis of Right ventricular Myocardial infarction in the presence of Inferior wall Myocardial infarction.

METHODS

Study Subjects:

Included in this prospective study were 50 consecutive patients (46 men and 4 women; mean [\pm SD] age, 56 ± 8 years) with inferior MIs who had been admitted to the coronary care unit Government Stanley medical college chennai at the onset of acute chest pain within the previous 24 h during period from 2007 to 2008. A standard 12-lead ECG and a right chest ECG (V_4 through V_6 R leads) were recorded Chest pain lasting for > 30 min, characteristic ST-segment elevation of ≥ 0.1 mV in two or more inferior derivation (i.e, leads II, III, and a VF) on ECGs, and a creatine kinase MB value more than twice that of the highest reference laboratory value were defined as acute MI criteria. The localization also was confirmed by the detection of hypokinetic and /or akinetic segments by echocardiography. The presence of an RVMI in association with an

inferior MI was defined by an ST-segment elevation of > 0.1 mV in V_4 through V_6 R lead derivations. Accordingly, RVMI was identified in 15 patients. Thirty six patients received thrombolytic therapy.

Exclusion criteria

Patients who had experienced MIs in the past, or had valvular disease, left and right bundle-branch block, hypertension and diabetes-atrial fibrillation were excluded from the study.

ECHOCARDIOGRAPHY

Two-dimensional, pulsed Doppler and color-flow Doppler echocardiography examinations were performed within two days after the onset of symptoms. A cardiac ultrasonographic unit (Aloka Prosound 4000) all equipped with a variable-frequency phased-array transducer and RD capabilities was used. All the echocardiographic measurements were performed according to the recommendations of the American Echocardiography Association without any information about the ECG findings.

ANNULAR VELOCITIES OBTAINED BY TDI

Pulsed-wave TDI images were acquired by activating the TDI function of the cardiac ultrasonographic unit. The best recordings were

obtained by optimizing the gains to terminate the signals formed by the transmittal flow and to reduce the noise with the use of low wall filter settings (ie, 50Hz). A 3.5-mm sample volume was used. From the apical four-chamber view, the TDI cursor was placed to the right of the ventricular free wall and the interventricular septum at the level of the tricuspid annulus in such a way that the annulus moved along the sample volume line.

A major positive S_m was recorded with the movement of the annulus towards the cardiac apex during systole. Two major negative velocities were recorded with the movement of the annulus toward the base of the heart during diastole, as follows: one during the early phase of diastolic myocardial velocity (E_m) velocity and another during the late phase of diastolic myocardial velocity (A_m). In the TDI images, S_m duration was measured as the ejection time (ET), the time between the end of the S_m and the beginning of the E_m as isovolumetric relaxation time (IVRT), and the time between the end of A_m and the beginning of S_m as isovolumetric contraction time (IVCT)

The right ventricular MPI was calculated as $(IVRT + IVCT)/ET$, by using the values obtained from the right ventricular free wall. In this study, Doppler velocity range of -20 to 20 cm/s was selected, and the velocities were measured on-line at a sweep of 50 mm/s. A mean of three consecutive cycles was used to calculate all Doppler echocardiographic

parameters.

CORONARY ANGIOGRAPHY

Coronary angiography were performed to detect the IRA within the first month after the MI. Total or subtotal occlusion of the coronary artery supplying the asynergic field was accepted as the defining features for the IRA. The patients were divided into two groups according to the level of the right coronary artery lesion when the IRA was the right coronary artery. The patients with the lesion proximal to the marginal branch of the right coronary artery were defined as group I, where those with the lesion distal to the marginal branch of right coronary artery were defined as groups II. The patients in whom the IRA is the circumflex artery were defined as group III. Those patients in whom the IRA could not be identified were excluded from the study. Coronary angiograms using the Judkins technique were performed with a standard cineangiographic system in multiple views. Lumen diameter narrowing was graded as 0,25,50,70,90 and 100%. In the present study, the definition of a significant anatomical coronary artery stenosis implied a $> 70\%$ localized luminal narrowing.

STATISTICAL ANALYSIS:

Variable are presented as mean \pm 1 SD. Analysis of variance was performed to estimate intergroup differences. Linear regression analyses and partial correlation testing using Pearson's method were used to assess

univariate relations. The prediction of RVMi was made using stepwise, forward, multiple regression analyses that included potential confounding variables not obviously related to each other. The null hypothesis was rejected for $p < 0.05$.

RESULTS

The clinical parameters were comparable in the groups based on the IRA. The number of the patients receiving thrombolytic therapy and the values of creatine kinase and transaminase did not differ between the groups. The percentage of the occlusion of the IRA were co

RESULTS

The clinical parameters were comparable in the groups based on the IRA. The number of the patients receiving thrombolytic therapy, and the peak values of creatine kinase and transaminase did not differ between the groups. The percentage of the occlusion of the IRA were comparable. (Table 1).

TABLE – 1

Comparison of the Baseline Clinical and Angiographic Parameters Among Patient Groups

Variables	Groups I (n = 23)	Group II (n = 17)	Group III (n = 10)
Age, yr	53 \pm 9	56 \pm 7	57 \pm 6
SBP, mm Hg	108 \pm 12	102 \pm 26	114 \pm 10
DBP, mm Hg	70 \pm 8	72 \pm 10	74 \pm 10
Heart rate, beats / min	68 \pm 12	70 \pm 10	76 \pm 8
BMI	26 \pm 3	28 \pm 4	26 \pm 3
Thrombolytic Therapy	20	10	6
Creatine Kinase MB	116 \pm 40	108 \pm 15	150 \pm 21
SGOT	123 \pm 30	116 \pm 20	140 \pm 35
IRA lesion, %	92 \pm 6	86 \pm 11	90 \pm 8

* Values given as mean \pm SD. SBP = Systolic BP; DBP = Diastolic BP; BMI = Body Mass Index; SGOT = Serum Glutamate Oxalate Transaminase; IRA = Infarct Related Artery.

Annulus Velocities and MPI Obtained by TDI:

In the comparison of patient groups with and without RVMI (based on the IRA), the values for Sm, Em, and Am obtained from the interventricular septum were lower in those with RVMIs than in those without, but the difference was insignificant. However, the Sm, Em and ET values obtained from the right ventricular free wall were significantly lower, and IRT and ICT values were significantly higher in those with RVMIs than in those without. The right ventricular MPI was calculated to be higher in those with RVMI than in those without RVMI.

TABLE - 2

Comparison of the Tissue Doppler Parameters Obtained from the Interventricular Septum in Patients With and Without RVMI

Parameters	RVMI (+) (n = 15)	RVMI (-) (n = 35)
Interventricular septum		
Sm, cm/s	5.8 \pm 1.3	6.6 \pm 1.5
Em, cm/s	5.3 \pm 1.4	5.6 \pm 1.7
Am, cm/s	7.6 \pm 2.2	8.8 \pm 2.3
Em / Am	0.76 \pm 0.31	0.68 \pm 0.25
IRT, ms	111 \pm 12	116 \pm 21
ICT, ms	74 \pm 12	82 \pm 18
ET, ms	258 \pm 24	250 \pm 12

Values given as mean \pm SD. P value – Non Significant

TABLE - 3

Comparison of the Tissue Doppler Parameters Obtained from the Right ventricular
free wall in Patients With and Without RVMI

Parameters	RVMI (+) (n = 15)	RVMI (-) (n = 35)
Right ventricular freewall		
Sm, cm/s	10.8±1.2	14.2±3.1
Em, cm/s	8.5±1.7	10.0±3.3
Am, cm/s	13.7±3.4	14.7 ±4.3
Em / Am	0.64±0.15	0.72±0.24
IRT, ms	114 ±12	88 ±8
ICT, ms	82 ±16	70±15
ET, ms	240±15	280 ± 22
RV-MPI	0.83±0.12	0.57±0.11

Values given as mean ± SD. P value – Non Significant

TABLE - 4

Comparison of the Right Ventricular Velocities and the MPI by Patient Group

Variables	Group I (n-23)	Group II (n-17)	Group III (n-10)	P Valve
Sm, cm/s	11.3 \pm 2.5	15.0 \pm 2.9	14.6 \pm 2.58	P< 0.01 (I compared with ground II & III).
Em. cm/s	8.6 \pm 2.6	9.3 \pm 1.8	12.2 \pm 4.1	P<0.01 (I compared with ground III).
Am, cm/s	13.6 \pm 3.7	14.7 \pm 4.5	15.8 \pm 4.5	NS
Em / Am	0.65 \pm 0.17	0.67 \pm 0.21	0.80 \pm 0.31	NS
IRT, ms	107 \pm 18	83 \pm 17	88 \pm 15	P<0.001 (I compared with group II).
ICT, ms	77 \pm 13	74 \pm 18	63 \pm 14	P<0.005 (I compared with group III).
ET, ms	252 \pm 20	286 \pm 26	284 \pm 16	P<0.001 (I compared with group II & III).
MPI	0.74 \pm 0.13	0.56 \pm 0.15	0.54 \pm 0.07	P<0.001 (I compared with group II & III).

Values given as mean \pm SD. NS = Not Significant.

The Role of Right Ventricular Sm and MPI Obtained by TDI in the Identification of RVMI and the Localization of IRA:

In the patient groups, the right ventricular free wall mean Sm was 10.8 ± 1.2 cm/s in group I and the mean right ventricular MPI was 0.74 ± 0.13 . Accordingly, Sm < 12 cm/s identifies RVMI and IRA (the proximal right coronary artery lesion) with the sensitivity (76%), Specificity (93%), negative predictive value (84%), and positive predictive value (86%), MPI of >0.70 identifies RVMI and IRA (the proximal right coronary artery lesion) with the sensitivity (88%), Specificity (81%) negative predictive value (96%), and positive predictive value (72%).

DISCUSSION

Right ventricular function is the prognostic predictor of numerous heart diseases, and it is affected especially by RVMI that accompanies inferior MI resulting from the occlusion of the right coronary artery. RVMI impairs right ventricular function. However, it is difficult to assess both right ventricular anatomy and function with reliable conventional echocardiographic examination because of its complex structure. In most cases, the visual assessment of the right ventricular free wall by echocardiography leads to an underestimation of hypokinesia, due to an asymmetric contraction of the right ventricular walls toward its center. It has been suggested that color or pulsed-wave TDI, which is frequently used in assessing left ventricular function, also can be used to assess right ventricular function. The myocardial velocities obtained by pulsed-wave TDI from the right ventricular free wall at the level of the tricuspid annulus in the apical four-chamber view show the longitudinal dynamics of the right ventricle, which is not seen with visual scoring.

In studies assessing right ventricular function by using techniques other than TDI, the movement of the tricuspid annulus has been shown to represent the global right ventricular function. It has been documented that significant right coronary artery disease can be identified with the assessment of systolic velocity obtained by pulsed-wave TDI from the right ventricular free wall close to the tricuspid lateral annulus in the apical four-chamber view during dobutamine stress echocardiography. Finally, it has been shown that tricuspid annulus systolic velocity decreases in patients with inferior MIs combined with RVMI compared to those without RVMI. On the basis of the results of these studies, we studied the diagnostic value of the right ventricular free wall S_m in the identification of RVMI proximal right coronary artery disease. We found that the right ventricular free wall S_m and E_m obtained from the right ventricular free wall was significantly lower in the patients in whom the IRA was the proximal right coronary artery than in those in whom the IRA was the distal right or circumflex coronary artery.

The decreased S_m of the right ventricular free wall can be accepted as an indicator of right ventricular systolic dysfunction, consequent to RVMI. Also, a decrease in the E_m of the right ventricular free wall shows the development of right ventricular diastolic dysfunction in those with RVMI and in those who experienced MIs due to a proximal right coronary artery lesion. An S_m of < 12 cm/s showed RVMI with high sensitivity (76%) and specificity (93%). In addition, the negative predictive value was also high (84%).

The MPI was defined as a noninvasive Doppler measurement of ventricular function. The MPI, also known as the Tei index, is commonly used in the assessment of systolic and diastolic function of the left ventricle. It has been reported that the MPI obtained by the combination of systolic and diastolic time intervals

can be used in the determination of the degree of dysfunction in patients with left ventricular dysfunction, and it correlates with conventional parameters such as left ventricular ejection fraction and invasive measurements. It also has been shown that this index is a simple and useful method that is independent of heart rate and is unaffected by the geometric shape of the ventricle, with excellent reproducibility between observers. Tei et al have shown that the right ventricular MPI is increased (0.93) in patients with primary pulmonary hypertension, and it is the most important indicator in discriminating healthy subjects. Eidem et al also have shown that the MPI increases significantly (0.63) in subjects with severe right ventricular insufficiency and Epstein anomaly. We considered calculating the MPI with TDI. We detected that the MPI that was calculated with this method was significantly higher in patients with RVMI in whom the IRA was the proximal right coronary artery. A right ventricular MPI of > 0.70 correlates with identification of RVMI and proximal right coronary artery disease. In 14 of 15 patients with RVMI and in 21 of 23 patients in whom the IRA was the proximal right coronary artery, the MPI was > 0.70 , whereas in 24 of 35 patients without RVMI and in 23 of 27 patients in whom the IRA was the distal right or circumflex coronary artery it was ≤ 0.70 . These findings have demonstrated that an MPI of ≥ 0.70 may diagnose RVMI and proximal right coronary artery disease with high sensitivity and specificity.

The early diagnosis of RVMI still has been suboptimal. Although the clinical triad of hypotension, absence of pulmonary congestion, and elevated jugular venous pressure is quite specific, the sensitivity is only 10 to 25%. At the present time, even when using additional chest leads, electrocardiography, which is the most common modality used for the diagnosis of RVMI, remains insufficient when compared with autopsy-proven MI. RVMI resulting from the occlusion of the proximal right coronary artery leads to pandiastolic dysfunction as well as to right ventricular systolic dysfunction. The Tissue Doppler Imaging method assessing myocardial function impairment is a new diagnostic tool that can be used to detect the IRA localization in patients with inferior MIs. The right ventricular free wall Sm and MPI values obtained by this method can give correct and reliable information about the severity of disease and the localization of the lesion, as well as the diagnosis.

LIMITATIONS OF THE STUDY

It has been known that although coordinated right ventricular function depends on the contraction of myocardial fibers along both the long and short axes, myocardial velocities obtained by TDI in the apical four chambers reflect the movements of the myocardium only along the long axis. The contraction of ventricular circumferential fibers does not affect the myocardial velocities in the image in this localization. This can lead to the ideal that TDI does not reflect the global right ventricular function exactly.

Because imaging of the right ventricle is not easy due to technical difficulties, its assessment is not practical with echocardiography. Besides, it is accepted that mitral and tricuspid annulus changes are affected by the movement of the entire wall of the ventricle. That is why the change in the right ventricular free wall annulus was evaluated. As is known, the improvement in right ventricular function generally occurs early after MI.

Although echocardiographic assessment was performed within the first 2 days in this study, the possibility that some patients already had recovered from the damage in the right ventricle may decrease the diagnostic value of the parameters that were used.

However, the parameters obtained by TDI provide quantitative data, which can give an idea about the extent of the RVMI as well as diagnosing it. But a correlation of these data with autopsy findings that can be accepted as the only "gold standard" is needed.

CONCLUSION

- The right ventricular systolic velocity (tricuspid annular velocity) obtained from the free wall decreases and the right ventricular myocardial performance index increases in patients with Right ventricular myocardial infarction.
- The Right ventricular myocardial infarction in the presence of inferior wall MI could be diagnosed and a proximal right coronary artery lesion could be predicted correctly by the use of pulsed tissue doppler parameters, which are easily obtained by Tissue Doppler Imaging method.

BIBLIOGRAPHY

1. Cohn JN. Right Ventricular infarction revisited. *Am J Cardiol* 1979;43:666-668.
2. Isner JM. Right ventricular myocardial infarction. In: Konstam MA, Isner JM, eds. *The right ventricle*. Boston : Kluwer Academic, 1988:87-129.
3. Cohn JN, Guha NH, Broder MI, Limas CJ. Right ventricular infarction : clinical and hemodynamic features. *Am J Cardiol* 1974;33:209-214.
4. Berger PB, Ryan TJ. Inferior myocardial infarction : high-risk subgroups. *circulation* 1990;81:401-411.
5. Bates ER, Elemmensen PM, Califf RM, et al. Precordial ST segment depression predicts a worse prognosis in inferior infarction despite reperfusion therapy. *J Am Coll Cardiol* 1990;16:1538-1544.
6. Nasmith J, Marpole D, Rahal D, Homan J, Stewart S, Sniderman A. Clinical outcomes after inferior myocardial infarction. *Ann Intern Med*. 1982;96:22-26.
7. Forman MB, Goodin J, Phelan B : Electrocardiographic changes associates with isolated right ventricular infarction. *J Am Coll Cardiol* 1984 Sep; 4(#); 640-3.

8. Mittal SR : Isolated right ventricular infarction. *Int. J Cardiol* 1994 Aug; 46(1) : 53-60.
9. Marvic Z, Zaputovic L, Matana A : Prognostic significance of complete atrioventricular block in patients with acute inferior myocardial infarction with and without right ventricular involvement. *Am Heart J* 1990 Apr; 119(4) : 823-8.
10. Robalino BD, Petrella RW, Jubran FY : Atrial natriuretic factor in patients with right ventricular infarction. *J Am Coll Cardiol* 1990 Mar 1; 15(3) : 546-53.
11. Robalino BD, Whitlow PL, Underwood Discipline Authority, Salcedo EE. Electrocardiographic manifestations of right ventricular infarction. *Am Heart J* 1989; 118:138-144.
12. Bar FW, Vermeer F, De Zwaan C, et al. Value of admission electrocardiogram in predicting outcome of thrombolytic therapy in acute myocardial infarction : a randomized trial conducted by the Netherlands Interuniversity Cardiology Institute. *Am J Cardiol* 1987;59:6-
13. Strauss HD, Sobel BE, Roberts R : The influence of occult right ventricular infarction on enzymatically estimated infarct size, hemodynamics and prognosis. *Circulation* 1980 Sep ; 62(3) : 5038.

14. Singhal AM, Ilangovan S, Mehta S : Isolated right ventricular infarction followed by posterior left ventricular infarction after a few days. *Acta Cardiol* 1984; 39(4) : 307-12.
15. Sugimoto T, Ogawa K, Asada T : Surgical treatment of ventricular septal perforation with right ventricular infarction. *J Cardiovasc Sug. (Torino)* 1996 Feb; 37(1):71-4.
16. Sugiura T, Iwasaka T, Shiomi K : Clinical significance of right ventricular dilatation in patients with right ventricular infarction. *Coron Artery Dis* 1994 Dec; 5(12) : 955-9.
17. Greenbaum RA, Ho SY, Gibson DG, Becker AF, Anderson RH. Left ventricular fibre architecture in man. *Br Heart J* 1984; 45:248-263.
18. Hamilton WF, Romf JH, Movements of the base of the ventricle and the relative constancy of the cardiac volume. *Am J Physiol* 1932;104:559-565.
19. Feigenbaum H, Zaky A, Nasser WK. Use of ultrasound to measure left ventricular stroke volume. *Circulation* 1967;38:1092-1099.
20. Simonson J, Schiller NB. Descent of the base of the left ventricle : an echocardiographic index of left ventricular function. *J Am Soc Echocardiogram* 1989;2:25-35.
21. Jones CJH, Raposo L, Gibson DG. Functional importance of the long axis dynamics of the human left ventricle. *Br Heart J* 1990;63:215-220.

22. Pai RG, Bodenheimer MM, Pai SM, Koss JH, Adamick RD.
Usefulness of systolic excursion of the mitral annulus as an index of left ventricular systolic function. *Am J Cardiol* 1991;67:222-224.
23. Gulati VK, Katz WE, Follansbee WP, Gorscan J. Mitral annular descent velocity by tissue Doppler echocardiography as an index of global left ventricular function. *Am J Cardiol* 1996;77:979-984.
24. Willenheimer R, Cline C, Erhardt L, Israelsson B. Left ventricular atrio-ventricular plane displacement : An echocardiographic technique for rapid assessment of prognosis in heart failure. *Heart* 1997;78:230-236.
25. Rodriguez L, Garcia M, Ares M, Griffin BP, Nakatani S, Thomas JD, Assessment of mitral annular dynamics during diastole by Doppler tissue imaging : comparison with mitral Doppler inflow in subjects without heart disease and in patients with left ventricular hypertrophy. *Am J Heart* 1996;131:982-987.
26. Oki T, Tabata T, Yamada H, Wakatsuki T, Shinohara H, Nishikado A, Luchi A, Fukuda N, Ito S. Clinical application of pulsed Doppler tissue imaging for assessing abnormal left ventricular relaxation. *Am J Cardiol* 1997;79:921-928.
27. Sohn DW, Chai IH, Lee DJ, Kim HC, Kim HS, Oh BH, Lee MM, Park YB, Choi YS, Seo JD, Lee YW. Assessment of mitral annulus velocity by Doppler tissue imaging in the evaluation

- of left ventricular diastolic function. J Am Coll Cardiol 1997;30:474-480.
28. Ramdas PG, Kanwaljit SG, Amplitudes, durations, and timings of apically directed left ventricular myocardial velocities : II. Systolic and diastolic asynchrony in patients with left ventricular hypertrophy. J Am Soc Echocardiogram 1998;11:112-118.
 29. Nagueh, SF, Sunn H, MD, Kopelen HA, Middleton KH, Khoury DS. Hemodynamic determinants of the mitral annulus diastolic velocities by tissue Doppler. J Am Coll Cardiol.
 30. San Roman, JA, Vilacosta, I, Rollan, MJ, et al Right ventricular asynergy during dobutamine-atropine echocardiography. J Am Coll Cardiol 1997;30, 430-435.
 31. Isaz, K, Munoz del Romeral, L, Lee, E, et al Quantitation of the motion of cardiac base in normal subjects by Doppler echocardiography. J Am Soc Echocardiogram 1993;6, 166-176.
 32. Kaul, S, Tei, C, Hopkins, JM, et al Assessment of right ventricular function using two-dimensional echocardiography. Am Heart J 1984;107,526-531.
 33. Wilson, NJ, Neutze, JM, Rutland, MD, et al Transthoracic echocardiography for right ventricular function late after the Mustard operation. Am Heart J 1996;131,360-367.

34. Rambalu, R, Poldermans, D, Fioretti, PM, et al Usefulness of pulse-wave Doppler tissue sampling a doubutamine stress echocardiography for the diagnosis of right coronary artery narrowing. *Am J Cardiol* 1998;81,1411-1415.
35. Kinch, JM, Ryan, TJ (1994) Right ventricular infarction. *N Engl J Med* 330, 1211-1217.
36. Berger, PB, Ryan, TJ Inferior myocardial infarction : high risk subgroups. *Circulation* 1990;81,401-411.
37. San Roman, JA, Vilacosta, I, Rollan, MJ, et al Right ventricular asynergy during doubutamine-atropine echocardiography. *J Am Coll Cardiol* 1997;30,430-435.
38. Isaz, K, Munoz del Romeral, L, Lee, E, et al Quantitation of the motion of cardiac base in normal subjects by Doppler echocardiography. *J Am Soc Echocardiogr* 1993;6,166-176.
39. Kaul, S, Tei, C, Hopkins, JM, et al Assessment of right ventricular function using two-dimensional echocardiography. *Am Heart J* 1984;107,526-531.
40. Wilson, NJ, Neutze, JM, Rutland, MD, et al Trasnthoracic echocardiography for right ventricular function late after the Mustard operation. *Am Heart J* 1996;131, 360-367.
41. Rambaldi, R, Poldermans, D, Fioretti, PM, et al Usefulness of pulse-wave Doppler tissue sampling and doubutamine stress

echocardiography for the diagnosis of right coronary artery narrowing. *Am J Cardiol* 1998;81, 1411-1415.

42. Schiller, NB, Shah, PM, Crawford, M, et al Recommendations for quantitation of the left ventricle by 2-dimensional echocardiography. *J Am Soc Echocardiogr* 1989;2, 358-367.
43. Currie, PJ, Seward, JB, Chan, KL, et al Continuous wave Doppler determination of right ventricular pressure: a simultaneous Doppler-catheterization study in 127 patients. *J Am Coll Cardiol* 1985;6, 750-756.
44. Nagueh, SF, Middleton, KJ, Kopelen, HA, et al Doppler tissue imaging: a new non-invasive technique for evaluation of left ventricular relaxation and estimation of filling pressures. *J Am Coll Cardiol* 1997;30, 1527-45.
45. Eidem, BW, O'Leary, PW, Tei, C, et al Usefulness of the myocardial performance index for assessing right ventricular function in congenital heart disease. *Am J Cardiol* 2000;86,654-658.

ABBREVIATIONS

Am = late phase of diastolic myocardial velocity;

E/A = E-wave/A-wave ratio;

Em = early phase of diastolic myocardial velocity;

ET = ejection time;

IVCT = isovolumetric contraction time ;

IRA = infarct-related artery;

IVRT = isovolumetric relaxation time;

MI = myocardial infarction;

MPI = myocardial performance index;

PD = pulmonary peak diastolic velocity;

PS = venous peak systolic velocity;

RVMI = right ventricular myocardial infarction;

Sm = myocardial systolic velocity;

TDI = tissue Doppler imaging;

PROFORMA

TISSUE DOPPLER IMAGING FOR THE DIAGNOSIS OF RIGHT VENTRICULAR MYOCARDIAL INFARCTION AND IDENTIFICATION OF INFARCT RELATED ARTERY IN ACUTE INFERIOR WALL MYOCARDIAL INFARCTION

Name	Age	Sex
BP	Heart rate	BMI
DM	HT	Smoking
Alcoholic	CKMB	SGOT

ECG

Thrombolytic therapy : YES/ NO

ECHO 2D

TDI : RV FREE WALL

SEPTAL WALL

Sm(cm/s)	IRT(ms)	Sm(cm/s)	IRT(ms)
Em(cm/s)	ICT(ms)	Em(cm/s)	ICT(ms)
Am(cm/s)	ET(ms)	Am(cm/s)	ET(ms)
Em/Am	MPI	Em/Am	

Coronary angiogram : Lesion site

Percentage :